

towards lower angle.

Upper side of hind wing the same.

Under side of fore wing dark brown with golden lustre. Basal area dusted with lustrous green scales as on upper surface. Inner marginal area very light brown.

Under side of hind wing the same as upper surface. Taken in January, 1900. This is practically identical with *T. pervivax*, Hubn. excepting that *pervivax* has a very prominent semitransparent band crossing the fore wing. The specimens in hand show no indication of such a band except in one instance where I noticed a slight dot. It would be unsafe to create a new species and the advisability of naming a variety is doubtful. There is a specimen in the Godman collection un-named.

*Thanaos parkeri* sp. nov.

Habitat: Suapure, Venezuela.

Expanse: 1.60 inches.

Head, thorax and abdomen above, very dark brown. Antennae black with white annulations at base of each joint. Club black above; tawny brown beneath, and approaching white towards base. Legs nearly black.

Upper side of fore wing very dark brown with slight grayish dustings. Near the base is a slight suggestion of gray dusting running from costa towards inner margin scarcely perceptible. The interspaces beyond, in centre of wing and the centre of discoidal space show a slight gray dusting. Outside of this in a line from costa downwards to lower submedian nervule, is a slightly prominent irregular band of gray dusting, broad at costa but diminishing and disappearing towards lower submedian nervule. In apical area outside of this band are four subcostal semitransparent white dots. Hind margin edged with a dark thread, and the hind marginal area has a series of interspacial grayish dashes from costa down to inner margin, well within the margin, and hardly noticeable.

Upper side of hind wing dark brown. The only marking is a suggestion of light brown dashes in interspaces running from upper angle to a point just above anal angle on inner margin. Hind margin edged with a dark thread. Fringe is generous and near upper angle is edged with white.

Under side of fore wing dead dark brown. The gray dustings of upper

surface do not show. The series of interspacial dashes or dustings within the hind margin are fairly prominent. The subcostal white dots are repeated. The hind margin has a dark thread and the fringe shows light hairs in interspaces.

Under side of hind wing dead dark brown. The line of interspacial light dashes of upper surface, running from upper angle downward are repeated and more clearly defined.

This species is closely allied to *N. tatus* Edw., *N. juvenalis* Fabr. and that group, and a detailed and identifiable description is almost impossible. One specimen was found unnamed in the Godman collection; also one specimen unnamed in the British Museum, taken in Malto Grosso, Brazil.

Taken in December, 1900.

***Nisoniades plummertini* sp. nov.**

Habitat; Suapure, Venezuela.

Expanse .95 inch.

Head, antennae, thorax, abdomen and legs above, very dark brown; below, lighter brown.

Upper side of fore wing dark brown. Across centre of wing from centre of costa a darker band runs downwards through discoidal space to inner margin. A similar band, but slightly broader, starts from costa and runs down to inner margin at a point near lower angle, following the contour of hind margin. On the inner edge of this band are four subcostal white dots and two more below them in the first and second submedian interspaces. There is a very slight dot also at the end of the discoidal space. The interspaces at edge of hind margin are slightly darker than ground color.

Upper side of hind wing the same as fore wing, the bands being continued to inner margin. The dots are, however, wanting.

Under side of fore wing the same as upper surface except that color is lighter and deader.

Under side of hind wing the same as upper surface except that color is lighter and deader.

Taken in November, 1900.

A specimen was found in the Godman collection unnamed. It is very close to *N. hayhursti* Edw.